



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**Macewen, William,** *THE GROWTH AND SHEDDING OF THE ANTLER OF THE DEER. THE HISTOLOGICAL PHENOMENA AND THEIR RELATION TO THE GROWTH OF BONE.* Glasgow: Maclehose, Jackson & Co., publishers to the University. Pp. i-xvii, 1-109, with 109 half-tone figures. 1920.

This beautifully illustrated volume is of as great value to the anatomist as to any one interested in deer. For the first time we have an anatomical treatise on the growth and shedding of the deciduous antler of the deer. This important investigation deals in detail with the histological changes which take place in the growing antler; it adds much to our understanding of the process of shedding of the antlers and explains their phenomenal growth.

Nowhere among animals do we find such an enormous and rapid reproduction of all the elements of the skin as in the growing velvet of the deer, which may cover the largest palmate antler within three months. The blood vessels in the velvet, which also form in a surprisingly short time, maintain the temperature of the growing bone within the hairy covering. These vessels anastomose but little with those of the underlying bone. The osseous growth of the antler is so rapid that the different phases of bone development may occur coincidentally. From the very beginning of antler formation there is going on a preparation for shedding. The latter is caused by the constricting of the blood vessels within the antler, particularly at the base, through the rapid proliferation of the bony tissue around them, which is followed by necrosis. Prior to the shedding, the bone on the distal end of the pedicle becomes softened, blood vessels within the pedicle increase in number and size, and granulation tissue is formed which loosens the connection between the dead antler and the pedicle. The shedding of the velvet, which precedes that of the antler, is likewise provided for in early stages of its growth. The bony substance emanating from the pedicle overlaps the latter, even in the sprouting antler, and forms the corona. The corona sends at first bony projections between the blood vessels of the velvet, which thus lie well protected in grooves. Gradually, however, these vessels are compressed and ultimately strangled by the advancement of ossification in the corona. The whole cutis then dries and peels off. The abundant multiplication of the bony cells in the growing antler, which are derived from the osteoblasts in the pedicle, can hardly be explained by normal cell division alone. The process known as nuclear budding may also partly account for this rapid production in ossification. During the growing period the antler is capable of repair after injury; at this time the pedicle is full of blood, which it transmits to the interior of the antler. After the termination of its development, the antler, no doubt, dies. The pedicle, however, must at all times be in active life, and preparing for the production of a new antler immediately after the shedding of the old one.

In addition to these normal conditions, the effect of injury to and disease of the pedicle and of castration on the growth of antlers is discussed in this much needed book.

—A. H. Schultz.

**Krieg, H.** *UEBER DIE BILDUNG VON STREIFENZEICHNUNGEN BEI SÄUGETIEREN.* *Anatom. Anz.*, vol. 54, pp. 33-40. 1921.

A preliminary report of investigations on the causes and origin of the stripe-distribution in the mammalian skin. Three principal types of arrangement are